

Earlier abundance estimates

An abundance estimate of 29,610 common dolphins (CV=0.39) was obtained from an aerial survey program conducted from 1978 to 1982 on the continental shelf and shelf edge waters between Cape Hatteras, North Carolina and Nova Scotia (CETAP 1982). An abundance estimate of 22,215 (CV=0.40) common dolphins was obtained from a June and July 1991 shipboard line-transect sighting survey conducted primarily between the 200 and 2,000 m isobaths from Cape Hatteras to Georges Bank (Waring *et al.* 1992; Waring 1998). An abundance estimate of 1,645 (CV=0.47) common dolphins was obtained from a June and July 1993 shipboard line-transect sighting survey conducted principally between the 200 and 2,000 m isobaths from the southern edge of Georges Bank, across the Northeast Channel to the southeastern edge of the Scotian Shelf (NMFS 1993). An abundance estimate of 6,741 (CV=0.69) common dolphins was obtained from a July to September 1995 sighting survey conducted by two ships and an airplane that covered 32,600 km in waters from Virginia to the mouth of the Gulf of St. Lawrence. An abundance estimate of 30,768 (CV=0.32) common dolphins was generated from a line-transect sighting survey conducted during 6 July to 6 September 1998 by a ship and plane that surveyed 15,900 km of track line in waters north of Maryland (38°N)(Palka 2006). The SEFSC conducted a shipboard line-transect sighting survey between 8 July and 17 August 1998, surveying 4,163 km of track line in waters south of Maryland (38°N) and sighted no common dolphins (Mullin and Fulling 2003). Although the 1991, 1993, 1995, and 1998 surveys did not sample the same areas or encompass the entire common dolphin habitat (e.g., little effort in Scotian shelf edge waters), they did focus on segments of known or suspected high-use habitats off the northeastern USA coast. As recommended in the GAMMS Workshop Report (Wade and Angliss 1997), estimates older than eight years are deemed unreliable and should not be used for PBR determinations.

Recent surveys and abundance estimates

An abundance estimate of 6,460 (CV=0.74) common dolphins was obtained from an aerial survey conducted in July and August 2002 which covered 7,465 km of trackline over waters from the 1000 m depth contour on the southern edge of Georges Bank to Maine (Table 1; Palka 2006). The value of $g(0)$ used for this estimation was derived from the pooled data of 2002, 2004 and 2006 aerial survey data.

An abundance estimate of 90,547 (CV= 0.244) common dolphins was obtained from a line-transect sighting survey conducted during 12 June to 4 August 2004 by a ship and plane that surveyed 10,761 km of trackline in waters north of Maryland (38°N) (Table 1; Palka 2006.). Shipboard data were collected using the two independent team line transect method and analyzed using the modified direct duplicate method (Palka 1995) accounting for biases due to school size and other potential covariates, reactive movements (Palka and Hammond 2001), and $g(0)$, the probability of detecting a group on the track line. Aerial data were collected using the Hiby circle-back line transect method (Hiby 1999) and analyzed accounting for $g(0)$ and biases due to school size and other potential covariates (Palka 2005).

An abundance estimate of 30,196 (CV=0.537) common dolphins was derived from a shipboard survey of the U.S. Atlantic outer continental shelf and continental slope (water depths > 50m) between Florida and Maryland (27.5 and 38° N latitude) conducted during June-August, 2004 (Table 1). The survey employed two independent visual teams searching with 50x bigeye binoculars. Survey effort was stratified to include increased effort along the continental shelf break and Gulf Stream front in the Mid-Atlantic. The survey included 5,659 km of trackline, and accomplished a total of 473 cetacean sightings. Sightings were most frequent in waters north of Cape Hatteras, North Carolina along the shelf break. Data were corrected for visibility bias ($g(0)$) and group-size bias and analyzed using line-transect distance analysis (Palka, 1995, 2006; Buckland *et al.* 2001).

An abundance estimate of 84,000 (CV=0.36) common dolphins was obtained from an aerial survey conducted in August 2006 which covered 10,676 km of trackline in the region from the 2000 m depth contour on the southern edge of Georges Bank to the upper Bay of Fundy and to the entrance of the Gulf of St. Lawrence. (Table 1; Palka pers. comm.)

The findings and conclusions in this report have not been formally disseminated by NOAA and should not be construed to represent any agency determination or policy.

Table 1. Summary of abundance estimates for western North Atlantic common dolphin. Month, year, and area covered during each abundance survey, and resulting abundance estimate (N_{best}) and coefficient of variation (CV).			
Month/Year	Area	N_{best}	CV
Aug 2002	S. Gulf of Maine to Maine	6,460	0.74
Jun-Aug 2004	Maryland to Bay of Fundy	90,547	0.24
Jun-Aug 2004	Florida to Maryland	30,196	0.54
Jun-Aug 2004	Florida to Bay of Fundy (COMBINED)	120,743	0.23
Aug 2006	S. Gulf of Maine to upper Bay of Fundy to Gulf of St. Lawrence	84,000	0.36

Minimum Population Estimate

The minimum population estimate is the lower limit of the two-tailed 60% confidence interval of the log-normally distributed best abundance estimate. This is equivalent to the 20th percentile of the log-normal distribution as specified by Wade and Angliss (1997). The best estimate of abundance for common dolphins is 120,743 animals (CV =0.23) derived from the 2004 surveys. The minimum population estimate for the western North Atlantic common dolphin is 99,975.

Current Population Trend

A trend analysis has not been conducted for this species.

CURRENT AND MAXIMUM NET PRODUCTIVITY RATES

Current and maximum net productivity rates are unknown for this stock. For purposes of this assessment, the maximum net productivity rate was assumed to be 0.04. This value is based on theoretical modeling showing that cetacean populations may not grow at rates much greater than 4% given the constraints of their reproductive life history (Barlow *et al.* 1995).

POTENTIAL BIOLOGICAL REMOVAL

Potential Biological Removal (PBR) is the product of minimum population size, one-half the maximum productivity rate, and a “recovery” factor (MMPA Sec. 3. 16 U.S.C. 1362; Wade and Angliss 1997). The minimum population size is 99,975 animals. The maximum productivity rate is 0.04, the default value for cetaceans. The “recovery” factor, which accounts for endangered, depleted, threatened, or stocks of unknown status, relative to optimum sustainable population (OSP) is assumed to be 0.5 because the CV of the average mortality estimate is less than 0.3 (Wade and Angliss 1997). PBR for the western North Atlantic stock of common dolphin is 1,000.

ANNUAL HUMAN-CAUSED MORTALITY AND SERIOUS INJURY

Total annual estimated average fishery-related mortality or serious injury to this stock during 2001-2005 was 151 (CV=.11) common dolphins (Table 2).

Fishery information

Detailed fishery information is reported in Appendix III.

Earlier Interactions

Prior to 1977, there was no documentation of marine mammal bycatch in distant-water fleet (DWF) activities off the northeast coast of the U.S. With implementation of the Magnuson-Stevens Fisheries Conservation and Management Act (MSFCMA), an observer program was established which recorded fishery data and information on incidental bycatch of marine mammals. During the period 1977-1986, observers recorded 123 mortalities in foreign *Loligo* squid-fishing activities. No mortalities were reported in foreign *Illlex* squid fishing operations.

The findings and conclusions in this report have not been formally disseminated by NOAA and should not be construed to represent any agency determination or policy.

From 1977 to 1991, observers recorded 110 mortalities in foreign mackerel-fishing operations (Waring *et al.* 1990; NMFS unpublished data). This total includes one documented take by a U.S. vessel involved in joint-venture fishing operations. A U.S. joint venture (JV) mackerel fishery was conducted in the mid-Atlantic region from February-May 1998. Seventeen incidental takes of common dolphin were observed in this fishery.

In the Atlantic pelagic longline fishery between 1990 and 2005, 20 common dolphins were observed hooked and released alive.

Eight hundred and sixty-one common dolphin mortalities were observed between 1989 and 1998 in the pelagic drift gillnet fishery, resulting in an estimated annual mortality and serious injury attributable to this fishery of (CV in parentheses) 540 in 1989 (0.19), 893 in 1990 (0.18), 223 in 1991 (0.12), 227 in 1992 (0.09), 238 in 1993 (0.08), 163 in 1994 (0.02), 83 in 1995 (0), 106 in 1996 (0.07) and 255 in 1998 (0).

Twelve mortalities were observed in the pelagic pair trawl between 1991 and 1995. The estimated annual fishery-related mortality and serious injury attributable to this fishery (CV in parentheses) was 5.6 in 1991 (0.53), 32 in 1992 (0.48), 35 in 1993 (0.43), 0 in 1994 and 5.6 in 1995 (0.35).

The estimated fishery-related mortality of common dolphins attributable to the *Loligo* squid portion of the Southern New England/Mid-Atlantic Squid, Mackerel, Butterfish Trawl fisheries was 0 between 1997-1998 and 49 in 1999 (CV=0.97). After 1999 this fishery is included as a component of the mid-Atlantic bottom trawl fishery.

In the Atlantic mackerel portion of the Southern New England/Mid-Atlantic Squid, Mackerel, Butterfish Trawl fisheries, the estimated fishery-related mortality was 161 (CV=0.49) animals in 1997 and 0 in 1998 and 1999. However, the estimates in both the mackerel and *Loligo* fisheries should be viewed with caution due to the extremely low (<1%) observer coverage. After 1999 this fishery is included as a component of the mid-Atlantic bottom trawl and mid-Atlantic mid-water trawl fisheries.

There was one observed take in the Southern New England/mid-Atlantic Bottom Trawl fishery reported in 1997. The estimated fishery-related mortality for common dolphins attributable to this fishery was 93 (CV= 1.06) in 1997 and 0 in 1998 and 1999. After 1999 this fishery is included as a component of the mid-Atlantic bottom trawl fishery.

No common dolphins were taken in observed mid-Atlantic gillnet fishery trips during 1993 and 1994. Two common dolphins were observed taken in 1995, 1996 and 1997, and no takes were observed from 1998 to 2004. Using the observed takes, the estimated annual mortality (CV in parentheses) attributed to this fishery was 7.4 in 1995 (0.69), 43 in 1996 (0.79), 16 in 1997 (0.53), and 0 in 1998-2004.

For more details on the earlier fishery interactions see Waring *et al.* 2007.

Northeast Sink Gillnet

Four common dolphins were observed taken in northeast sink gillnet fisheries in 2005. The estimated annual fishery-related mortality and serious injury attributable to the northeast sink gillnet fishery (CV in parentheses) was 0 in 1995, 63 in 1996 (1.39), 0 in 1997, 0 in 1998, 146 in 1999 (0.97), 0 in 2000-2004 and 5 (0.80) in 2005. This fishery, which extends from North Carolina to New York, is actually a combination of small vessel fisheries that target a variety of fish species, some of which operate right off the beach. The number of vessels in this fishery is unknown, because records which are held by both state and federal agencies have not been centralized and standardized.

Northeast Bottom Trawl

This fishery is active in New England waters in all seasons. One common dolphin was observed taken in 2002, three in 2004, and five in 2005 (Table 2). The estimated annual fishery-related mortality and serious injury attributable to the northeast bottom trawl fishery (CV in parentheses) was 27 in 2000 (0.29), 30 (0.30) in 2001, 26 (0.29) in 2002, 26 (0.29) in 2003, 26 (0.29) in 2004, and 32 (0.28) in 2005. The 2001-2005 average mortality attributed to the northeast bottom trawl was 28 animals (CV=0.13).

Mid-Atlantic Bottom Trawl

Three common dolphins were observed taken in mid-Atlantic bottom trawl fisheries in 2000, two in 2001, nine in 2004, and 15 in 2005 (Table 2). The estimated annual fishery-related mortality and serious injury attributable to the northeast bottom trawl fishery (CV in parentheses) was 93 in 2000 (0.26), 103 (0.27) in 2001, 87 (0.27) in 2002, 99 (0.28) in 2003, 159 (0.30) in 2004, and 141 (0.29) in 2005. The 2001-2005 average mortality attributed to the mid-Atlantic bottom trawl was 118 animals (CV=0.13).

The findings and conclusions in this report have not been formally disseminated by NOAA and should not be construed to represent any agency determination or policy.

Table 2. Summary of the incidental mortality of common dolphins (*Delphinus delphis*) by commercial fishery including the years sampled (Years), the number of vessels active within the fishery (Vessels), the type of data used (Data Type), the annual observer coverage (Observer Coverage), the mortalities recorded by on-board observers (Observed Mortality), the estimated annual mortality (Estimated Mortality), the estimated CV of the annual mortality (Estimated CVs) and the mean annual mortality (CV in parentheses).

Fishery ^a	Years	Vessels	Data Type ^b	Observer Coverage ^c	Observed Serious Injury	Observed Mortality ^d	Estimated Serious Injury	Estimated Mortality	Estimated Combined Mortality	Estimated CVs	Mean Annual Mortality
Northeast Sink Gillnet	01-05	unk	Obs. Data Weighout, Logbooks	.06, .04, .02, .03, .06, .07	0, 0, 0, 0, 0	0, 0, 0, 0, 4	0, 0, 0, 0, 0	0, 0, 0, 0, 26	0, 0, 0, 0, 26	0, 0, 0, 0, .8	5 (0.8)
Northeast Bottom Trawl ^e	01-05 ^e	unk	Obs. Data Dealer Data VTR Data	.01, .03, .04, .05, .12	0, 0, 0, 0, 0	0, 1, 0, 3, 5	0, 0, 0, 0	30, 26, 26, 26, 32	30, 26, 26, 26, 32	.30, .29, .29, .29, .28	28 (.13)
Mid-Atlantic Bottom Trawl ^e	01-05 ^e	unk	Obs. Data Dealer	.01, .01, .03, .03	0, 0, 0, 0, 0	2, 0, 0, 9, 15	0, 0, 0, 0	103, 87, 99, 159, 141	103, 87, 99, 159, 141	.27, .27, .28, .30, .20	118 (.13)
TOTAL											151 (.11)

- The fisheries listed in Table 2. reflect new definitions defined by the proposed List of Fisheries for 2005 (FR Vol. 69, No. 231, 2004). The 'North Atlantic bottom trawl' fishery is now referred to as the 'Northeast bottom trawl. The Illex, Loligo and Mackerel fisheries are now part of the 'mid-Atlantic bottom trawl' and 'mid-Atlantic midwater trawl' fisheries.
- Observer data (Obs. Data), used to measure bycatch rates, are collected within the Northeast Fisheries Observer Program. NEFSC collects landings data (Dealer reported data) which are used as a measure of total landings and mandatory Vessel Trip Reports (VTR) (Trip Logbook) that are used to determine the spatial distribution of landings and fishing effort.
- The observer coverages for the Northeast sink gillnet fishery are ratios based on tons of fish landed. North Atlantic bottom trawl and mid-Atlantic bottom trawl fishery coverages are ratios based on trips.
- The data used to predict bycatch rates to estimate annual mortality were pooled over the years 2001-2005. The data are treated as one data set and assumed to represent average fishing practices during the time period. Regression techniques within a model framework were applied to the pooled data set. Therefore, if there was no observed bycatch reported for any one given year, this does not imply that there was no bycatch during that year. The exception would be if year was selected by the model as an important factor associated with observing bycatch.
- A new method was used to develop preliminary estimates of mortality for the Mid-Atlantic and Northeast bottom trawl fisheries during 2000-2004. They are a product of bycatch rates predicted by covariates in a model framework and effort reported by commercial fishermen on mandatory vessel logbooks. This method differs from the previous method used to estimate mortality in these fisheries prior to 2000. Therefore, the estimates reported prior to 2000 can not be compared to estimates during 2000-2004.

CANADA

Between January 1993 and December 1994, 36 Spanish deep water trawlers, covering 74 fishing trips (4,726 fishing days and 14,211 sets), were observed in NAFO Fishing Area 3 (off the Grand Banks) (Lens 1997). A total of 47 incidental catches were recorded, which included 1 common dolphin. The incidental mortality rate for common dolphins was 0.007/set.

Other Mortality

From 2001 to 2005, 322 common dolphins were reported stranded between Maine and Florida (Table 3). The total includes mass stranded common dolphins in Massachusetts during 2002 (9 animals), 2004 (one event of 6 animals and one of 3 animals), and a total of 25 in 2005 in 4 separate events; and in North Carolina in 2001 (7 animals). Five of the 2005 Massachusetts stranded animals were released alive. One stranded common dolphin calf in New Jersey was relocated to a rehabilitation facility in 2005. In 2001, one stranding mortality in Virginia and another animal in North Carolina were designated as human interactions/fishing interactions. Similarly in 2002, one stranding in New York and another animal in Virginia were designated as human interactions/fishery interactions. Common dolphins were included in the UME (unusual mortality event) declared for Virginia in 2004 (Marine Mammal Commission 2005). The strandings were primarily bottlenose dolphins, but common dolphins were also involved. Human interactions were implicated in one of the 2004 Virginia common dolphin mortalities. In 2005, one stranding mortality in New York was designated as human interaction.

Four common dolphin strandings (6 individuals) were reported on Sable Island, Nova Scotia from 1996 to 1998 (Lucas and Hooker 1997; Lucas and Hooker 2000). One common dolphin was reported stranded in Halifax County, Nova Scotia in 2005 (Tonya Wimmer, pers. comm.).

Table 3. Common dolphin (*Delphinus delphis*) reported strandings along the U.S. Atlantic coast, 2001-2005.

STATE	2001	2002	2003	2004	2005	TOTALS
Maine	1	0	0	0	0	1
Massachusetts ^a	8	34	21	26	59	148
Rhode Island	0	1	2	1	0	4
Connecticut	0	0	0	0	0	0
New York	6	5	11	3	4	29
New Jersey	5	1	6	8	4	24
Delaware	1	1	1	2	1	6
Maryland	2	0	0	4	0	6
Virginia ^b	4	3	4	8	2	21
North Carolina ^d	14	0	62	4	1	81
Georgia	0	0	0	0	0	0
Florida	0	1	0	0	0	1
EZ	0	0	0	1	0	1
TOTALS	41	46	107	57	71	322

a. Massachusetts mass strandings (2002 - 9 animals; 2004 - 6 and 3; 2005 - 7,5,25, and 4).
b. Virginia reports 1 common dolphin found in a pound net in 2004.
c. Fishery Interactions (FI)/Human Interactions (HI) - North Carolina reported 1 HI, fishing gear, April 2001; Virginia - 1 FI March 2001).
d. North Carolina mass stranding (2001 - 7 animals).
e. 2002 FI, one in NY, one in Va.

The findings and conclusions in this report have not been formally disseminated by NOAA and should not be construed to represent any agency determination or policy.

Stranding data probably underestimate the extent of fishery-related mortality and serious injury because all of the marine mammals that die or are seriously injured may not wash ashore, nor will all of those that do wash ashore necessarily show signs of entanglement or other fishery-interaction. Finally, the level of technical expertise among stranding network personnel varies widely as does the ability to recognize signs of fishery interaction.

STATUS OF STOCK

The status of common dolphins, relative to OSP, in the U.S. Atlantic EEZ is unknown. The species is not listed as threatened or endangered under the Endangered Species Act. There are insufficient data to determine the population trends for this species. The total U.S. fishery-related mortality and serious injury for this stock is not less than 10% of the calculated PBR and, therefore, cannot be considered to be insignificant and approaching zero mortality and serious injury rate. The 2001-2005 average annual human-related mortality does not exceed PBR; therefore, this is not a strategic stock.

REFERENCES CITED

- Barlow, J., S.L. Swartz, T.C. Eagle, and P.R. Wade. 1995. U.S. Marine Mammal Stock Assessments: Guidelines for Preparation, Background, and a Summary of the 1995 Assessments. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-6, 73 pp.
- Buckland, S.T., D.R. Andersen, K.P. Burnham, and J.L. Laake. 1993. Distance sampling: Estimating abundance of biological populations. Chapman and Hall, New York, 446 pp.
- Buckland, S.T., D.R. Andersen, K.P. Burnham, J.L. Laake, D.L. Borchers, and L. Thomas. 2001. Introduction to Distance Sampling: Estimating abundance of biological populations. Oxford University Press, New York, 432 pp.
- CETAP. 1982. A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf. Cetacean and Turtle Assessment Program, University of Rhode Island. Final Report, Contract AA51-C78-48, Bureau of Land Management, Washington, DC, 538 pp.
- Cramer, J. 1994. Large pelagic logbook newsletter - 1993. NOAA Tech. Mem. NMFS-SEFSC-352. 19 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Evans, W.E. 1994. Common dolphin, white-bellied porpoise. Pp 191-224. *In*: S. H. Ridgway and R. Harrison (eds.). Handbook of marine mammals, Volume 5: The first book of dolphins. Academic Press, San Diego, CA.
- Gaskin, D.E. 1992. Status of common dolphin, *Delphinus delphis*, in Canada. *Can. Field-Nat.* 106:55-63.
- Gowans, S. and H. Whitehead. 1995. Distribution and habitat partitioning by small odontocetes in the Gully, a submarine canyon on the Scotian Shelf. *Can. J. Zool.* 73:1599-1608.
- Hain, J.H.W., R.K. Edell, H.E. Hays, S.K. Katona, and J.D. Roanowicz. 1981. General distribution of cetaceans in the continental shelf waters of the northeastern U.S. Pages II1-II277. *In*: CETAP (Cetacean and Turtle Assessment program), A characterization of marine mammals and turtles in the mid- and north Atlantic areas of the U.S. outer continental shelf, Annual Report for 1979. Contract No. AA551-CT8-48, U.S. Dept. of Interior, Bureau of Land Management, Washington, DC.
- Hamazaki, Toshihide. 2002. Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, North Carolina, USA to Nova Scotia, Canada). *Mar. Mammal Sci.* 18(4):920-939.
- Hiby, L. 1999. The objective identification of duplicate sightings in aerial survey for porpoise. Pages 179-189 *in*: G.W. Garner, S.C. Amstrup, J.L. Laake, B.F.J. Manly, L.L. McDonald, and D.G. Robertson. (eds). Marine Mammal Survey and Assessment Methods. Balkema, Rotterdam.
- Johnson, D.R., C.A. Brown, and C. Yeung. 1999. Estimates of marine mammal and marine turtle catch by the U.S. Atlantic pelagic longline fleet in 1992-1997. NOAA Technical Memorandum NMFS- SEFSC-418. 70 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Laake, J.L., S.T. Buckland, D.R. Anderson, and K.P. Burnham. 1993. DISTANCE user's guide, V2.0. Colorado Cooperative Fish & Wildlife Research Unit, Colorado State University, Ft. Collins, CO. 72 pp.
-

The findings and conclusions in this report have not been formally disseminated by NOAA and should not be construed to represent any agency determination or policy.

- Lens, S. 1997. Interactions between marine mammals and deep water trawlers in the NAFO regulatory area. ICES CM 1997/Q:8. 10 pp.
- Lucas, Z.N. and S.K. Hooker. 1997. Cetacean strandings on Sable Island, Nova Scotia, 1990-1996. Paper SC/49/06 presented to the IWC Scientific Committee, September 1997. 10 pp.
- Lucas, Z.N. and S.K. Hooker. 2000. Cetacean strandings on Sable Island, Nova Scotia, 1970-1998. Can. Field-Nat. 114:45-61.
- Marine Mammal Commission. 2006. US Marine Mammal Commission Annual Report to Congress, 2004. Marine Mammal Commission, Bethesda, MD. vi+163pp.
Available at: <http://www.nefsc.noaa.gov/nefsc/publication/> 70 pp
- Mullin, K.D. and G.L. Fulling. 2003. Abundance of cetaceans in the southern U.S. North Atlantic Ocean during summer 1998. Fish. Bull., U.S. 101:603-613.
- NMFS [National Marine Fisheries Service]. 1993. Cruise results, NOAA ship DELAWARE II, Cruise No. DEL 93-06, Marine mammal Survey. 5 pp. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026.
- Palka, D. 1995. Abundance estimate of the Gulf of Maine harbor porpoise. Rep. int. Whal. Commn (Special Issue) 16:27-50.
- Palka, D. 1996. Update on abundance of Gulf of Maine/Bay of Fundy harbor porpoises. U.S. Dep. Commer., Northeast Fish. Sci. Cent. Ref. Doc.96-04; 37 pp.
- Palka, D. and P.S. Hammond. 2001. Accounting for responsive movement in line transect estimates of abundance. Can. J. Fish. Aquat. Sci. 58:777-787.
- Palka, D. 2005. Aerial surveys in the northwest Atlantic: estimation of $g(0)$. In: Proceedings of the workshop on Estimation of $g(0)$ in line-transect surveys of cetaceans, ed. F. Thomsen, F. Ugarte, and P.G.H. Evans. ECS Newsletter No. 44 – Special Issue. April 2005. Pgs 12-7.
- Palka, D.L. 2006. Summer abundance estimates of cetaceans in US North Atlantic Navy Operating Areas. U.S. Dep. Commer., Northeast Fish. Sci. Cent. Ref. Doc. 06-03, 41 pp.
- Payne, P.M., L.A. Selzer, and A.R. Knowlton. 1984. Distribution and density of cetaceans, marine turtles, and seabirds in the shelf waters of the northeastern United States, June 1980-December 1983, based on shipboard observations. NOAA/NMFS Contract No. NA-81-FA-C-00023. 245 pp.
- Scott, G.P. and C.A. Brown. 1997. Estimates of marine mammal and turtle catch by the U.S. Atlantic pelagic longline fleet in 1994-1995. Miami Laboratory Contribution MIA-96/97-28.
- Selzer, L.A. and P.M. Payne. 1988. The distribution of white-sided (*Lagenorhynchus acutus*) and common dolphins (*Delphinus delphis*) vs. environmental features of the continental shelf of the northeastern United States. Mar. Mammal. Sci. 4(2):141-153.
- Sergeant, D. E., A. W. Mansfield, and B. Beck. 1970. Inshore records of cetacea for eastern Canada, 1949-68. J. Fish. Res. Bd. Can. 27:1903-1915.
- Wade, P.R. and R.P. Angliss. 1997. Guidelines for assessing marine mammal stocks: Report of the GAMMS Workshop April 3-5, 1996, Seattle, Washington. U.S. Dep. Commer., NOAA Tech. Memo. NMFS-OPR-12, 93 pp.
- Waring, G.T. 1998. Results of the summer 1991 R/V Chapman marine mammal sighting survey. U.S. Dep. Commer., Northeast Fish. Sci. Cent. Ref. Doc. 98-09, 21 pp.
- Waring, G.T., P. Gerrior, P.M. Payne, B.L. Parry, and J.R. Nicolas. 1990. Incidental take of marine mammals in foreign fishery activities off the northeast United States, 1977-1988. Fish. Bull., U.S. 88(2):347-360.
- Waring, G.T., C.P. Fairfield, C.M. Ruhsam, and M. Sano. 1992. Cetaceans associated with Gulf Stream features off the northeastern USA shelf. ICES Marine Mammals Comm. CM 1992/N:12, 29 pp.
- Waring, G.T., E. Josephson, C.P. Fairfield, and K. Maze-Foley. 2007. U.S. Atlantic and Gulf of Mexico marine mammal stock assessments – 2006. NOAA Tech. Memo. NMFS-NE-201, xxx pp.

The findings and conclusions in this report have not been formally disseminated by NOAA and should not be construed to represent any agency determination or policy.

- Westgate, A. J. 2005. Population structure and life history of short-beaked common dolphins (*Delphinus delphis*) in the North Atlantic. Ph.D thesis. Nicholas School of the Environment and Earth Sciences, Duke University, Beaufort, NC.
- Yeung, C. 1999a. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1998. NOAA Tech. Memo. NMFS-SEFSC-430, 26 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Yeung, C. 1999b. Revised Mortality Estimates of Marine Mammal Bycatch in 1992-1997 based on Serious Injury Guidelines. NOAA Tech. Memo. NMFS-SEFSC-429, 23 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Yeung, C., S. Epperly, and C.A. Brown. 2000. Preliminary revised estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet, 1992-1999. NMFS, Miami Lab. PRD Contribution Number 99/00-13. 58 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.
- Yeung, C. 2001. Estimates of marine mammal and marine turtle bycatch by the U.S. Atlantic pelagic longline fleet in 1999-2000. NOAA Tech. Memo. NMFS-SEFSC-467, 43 pp. Available from: NMFS, Southeast Fisheries Science Center, 75 Virginia Beach Dr., Miami, FL 33149.