

# **Northeast Multispecies Fishery**

## **Final Year-End Results for Fishing Year 2013**

- Tables 1 through 10: Total groundfish caught, landed, and discard estimates
- Tables 11 through 15: U.S./Canada Stocks Catch Evaluation
- Table 16: FY 2011 through FY 2013 GOM cod and haddock recreational catch evaluation

**Table 1: FY 2013 Northeast Multispecies Percent of Catch Limit Caught (%)**

Stock	Components with ACLs and sub-ACLs; (with AMs)								sub-components: No AMs	
	Total Groundfish	Groundfish Fishery	Sector*	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
GB Cod	84.8	87.0	86.8	101.0					46.0	42.8
GOM cod	96.5	104.9	90.2	48.9	131.5				34.7	5.7
GB Haddock	11.9	11.4	11.4	0.5		106.2			2.1	4.8
GOM Haddock	147.9	154.4	91.2	108.9	312.2	-			30.4	25.3
GB Yellowtail Flounder	44.7	36.1	36.5	0.4			90.3	63.7	NA	0.6
SNE Yellowtail Flounder	70.1	63.7	57.8	93.1			111.4		206.5	106.3
CC/GOM Yellowtail Flounder	86.7	79.4	80.9	31.5					130.2	271.3
Plaice	97.5	98.3	99.8	14.3					63.0	95.7
Witch Flounder	99.2	105.3	106.6	30.6					115.5	64.5
GB Winter Flounder	48.4	48.8	49.1	0.0					NA	36.5
GOM Winter Flounder	23.6	23.7	24.4	6.6					24.8	16.5
SNE/MA Winter Flounder	63.6	65.2	62.4	87.0					23.7	108.1
Redfish	38.5	39.5	39.6	10.9					17.3	1.8
White Hake	51.7	53.1	53.4	21.6					5.5	9.9
Pollock	47.1	38.1	38.1	40.2					104.9	103.7
Northern Windowpane	195.0	242.4	NA	NA					62.2	95.0
Southern Windowpane	105.3	113.7	NA	NA			70.5		67.9	146.4
Ocean Pout	26.9	16.9	NA	NA					62.6	116.4
Halibut	82.1	105.2	NA	NA					57.5	31.0
Wolffish	29.3	27.6	NA	NA					185.0	26.3

\*Sector sub-ACL values have been corrected from the report dated October 20, 2014 to include unrenewed permit potential sector contributions

Values in percent of live weight (%), "-" for values exactly equal to zero

Source: NMFS Greater Atlantic Regional Office  
October 29, 2014, run date of July 15, 2014

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting; (4) Observers and at-sea monitors via the Northeast Fisheries Observer Program. Differences with previous reports are due to corrections made to the database.

**Table 2: FY 2013 Northeast Multispecies Annual Catch Limits (mt)**

Stock	ACLs and sub-ACLs: With Accountability Measures (AMs)								sub-components: No AMs	
	Total ACL	Groundfish sub-ACL	Sector sub-ACL*	Common Pool sub-ACL	Recreational sub-ACL	Midwater Trawl Herring Fishery sub-ACL	Scallop Fishery sub-ACL	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
GB cod	1,907	1,807	1,776	32					20	80
GOM cod	1,470	1,316	812	18	486				103	51
GB Haddock	27,936	26,196	26,111	85		273			293	1,173
GOM Haddock	274	261	185	2	74	3			4	6
GB Yellowtail Flounder	208.5	154.5	152.6	1.9			41.5	4.0		4.3
SNE Yellowtail Flounder	665	585.7	487.5	98.2			43.6		7.0	28.0
CC/GOM Yellowtail Flounder	523	479	466	13					33	11
Plaice	1,482	1,420	1,395	25					31	31
Witch Flounder	751	610	599	11					23	117
GB Winter Flounder	3,641	3,528	3,506	22						113
GOM Winter Flounder	1,040	715	688	26					272	54
SNE/MA Winter Flounder	1,612	1,210	1,074	136					235	168
Redfish	10,462	10,132	10,092	40					110	220
White Hake	3,974	3,849	3,822	27					42	84
Pollock	14,921	12,893	12,802	91					936	1,092
Northern Windowpane	144	98	NA	98					2	44
Southern Windowpane	527	102	NA	102			183		55	186
Ocean Pout	220	197	NA	197					2	21
Halibut	96	52	NA	52					40	5
Wolffish	65	62	NA	62					1	3

\*Sector sub-ACL values have been corrected from the report dated October 20, 2014 to include unrenewed permit potential sector contributions

Values in metric tons of live weight

Source: NMFS Greater Atlantic Regional Office  
October 29, 2014

**Table 3: FY 2013 Northeast Multispecies Total Catch (mt)**

Stock	Total Catch	Groundfish Fishery	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery <sup>1</sup>	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
GB Cod	1,616.3	1,572.9	1,540.6	32.3					9.2	34.2
GOM cod	1,418.8	1,380.1	732.0	8.8	639.3				35.8	2.9
GB Haddock	3,330.1	2,977.5	2,977.1	0.4		290.0			6.1	56.5
GOM Haddock	405.7	402.9	169.2	2.2	231.5	0.0			1.3	1.6
GB Yellowtail Flounder	93.3	55.8	55.8	0.0			37.5	2.5	0.0	0.0
SNE/MA Yellowtail Flounder	466.1	373.3	281.9	91.4			48.6		14.5	29.8
CC/GOM Yellowtail Flounder	453.1	380.5	376.5	4.1					42.8	29.7
Plaice	1,444.6	1,395.2	1,391.6	3.6					19.6	29.8
Witch Flounder	745.2	642.3	638.9	3.4					27.1	75.8
GB Winter Flounder	1,763.1	1,722.0	1,722.0	0.0					0.0	41.0
GOM Winter Flounder	245.6	169.3	167.6	1.7					67.4	8.9
SNE/MA Winter Flounder	1,025.9	788.6	670.4	118.3					55.7	181.6
Redfish	4,023.5	4,000.6	3,996.2	4.4					19.0	3.9
White Hake	2,056.3	2,045.6	2,039.8	5.8					2.3	8.3
Pollock	7,029.1	4,915.0	4,878.4	36.5					981.7	1,132.4
Northern Windowpane	280.1	237.5	237.3	0.2					0.9	41.6
Southern Windowpane	554.7	115.9	86.0	30.0			129.1		37.3	272.4
Ocean Pout	59.3	33.2	27.3	5.9					1.5	24.6
Halibut	79.0	54.7	53.8	0.9					22.8	1.5
Wolffish	19.1	17.1	17.1	0.0					1.3	0.7

<sup>1</sup>Based on scallop fishing year March, 2013 through February, 2014

Values in metric tons of live weight

Sector and common pool include estimate of missing dealer reports

Any value for a non-allocated species may include landings of that stock or misreporting of species and/or stock area. These are northern windowpane, southern windowpane, ocean pout, halibut, and wolffish.

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting; (4) Observers and at-sea monitors via the Northeast Fisheries Observer Program. Differences with previous reports are due to corrections made to the database.

**Table 4: FY 2013 Northeast Multispecies Landings (mt)**

Stock	Total Landings	Groundfish Fishery	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
GB Cod	1,554.0	1,525.2	1,494.0	31.3					6.0	22.7
GOM cod	1,294.7	1,261.4	712.3	8.5	540.6				33.2	0.2
GB Haddock	2,969.9	2,696.4	2,696.0	0.4		272.7			0.2	0.5
GOM Haddock	382.6	382.0	148.3	2.2	231.5	0.0			0.6	0.0
GB Yellowtail Flounder	49.7	46.2	46.2	0.0			3.6	0.0	0.0	0.0
SNE/MA Yellowtail Flounder	376.5	361.5	271.0	90.5			1.5		8.3	5.3
CC/GOM Yellowtail Flounder	406.0	362.8	359.7	3.1					40.8	2.4
Plaice	1,301.5	1,290.3	1,287.1	3.1					10.4	0.8
Witch Flounder	629.4	602.5	599.6	2.9					24.7	2.2
GB Winter Flounder	1,717.5	1,716.8	1,716.8	0.0					0.0	0.7
GOM Winter Flounder	232.3	164.8	163.1	1.7					65.3	2.2
SNE/MA Winter Flounder	853.8	776.9	663.5	113.4					47.7	29.2
Redfish	3,621.1	3,614.0	3,610.6	3.4					6.5	0.6
White Hake	2,023.9	2,021.8	2,016.6	5.2					0.8	1.3
Pollock	5,517.2	4,808.9	4,773.0	35.8					258.2	450.2
Northern Windowpane	0.0	0.0	0.0	0.0					0.0	0.0
Southern Windowpane	22.9	0.1	0.0	0.1			0.0		20.8	2.0
Ocean Pout	0.0	0.0	0.0	0.0					0.0	0.0
Halibut	36.1	14.3	13.4	0.8					21.1	0.8
Wolffish	0.0	0.0	0.0	0.0					0.0	0.0

Values in metric tons of live weight

Sector and common pool include estimate of missing dealer reports

Any value for a non-allocated species may include landings of that stock or misreporting of species and/or stock area. These are northern windowpane, southern windowpane, ocean pout, halibut, and wolffish.

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting; (4) Observers and at-sea monitors via the Northeast Fisheries Observer Program. Differences with previous reports are due to corrections made to the database.

**Table 5: FY 2013 Northeast Multispecies Estimated Discards (mt)**

Stock	Total Discards	Groundfish Fishery	Sector	Common Pool	Recreational	Midwater Trawl Herring Fishery	Scallop Fishery	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
GB Cod	62.4	47.7	46.6	1.1					3.2	11.5
GOM cod	124.1	118.8	19.7	0.3	98.8				2.6	2.7
GB Haddock	360.2	281.1	281.1	0.0		17.2			5.9	56.0
GOM Haddock	23.1	20.8	20.8	0.0	0.0	0.0			0.7	1.6
GB Yellowtail Flounder	43.5	9.6	9.6	0.0			33.9	2.5	0.0	0.0
SNE/MA Yellowtail Flounder	89.5	11.8	10.9	0.9			47.1		6.1	24.5
CC/GOM Yellowtail Flounder	47.1	17.8	16.7	1.0					2.0	27.3
Plaice	143.1	105.0	104.5	0.4					9.2	28.9
Witch Flounder	115.8	39.8	39.3	0.5					2.4	73.6
GB Winter Flounder	45.6	5.3	5.3	0.0					0.0	40.3
GOM Winter Flounder	13.4	4.5	4.5	0.0					2.1	6.7
SNE/MA Winter Flounder	172.2	11.7	6.8	4.9					8.1	152.4
Redfish	402.4	386.6	385.6	0.9					12.5	3.3
White Hake	32.4	23.8	23.2	0.6					1.5	7.1
Pollock	1,511.9	106.1	105.4	0.7					723.6	682.2
Northern Windowpane	280.1	237.5	237.3	0.2					0.9	41.6
Southern Windowpane	531.8	115.9	86.0	29.9			129.1		16.5	270.4
Ocean Pout	59.3	33.2	27.3	5.9					1.5	24.6
Halibut	42.9	40.4	40.4	0.1					1.7	0.7
Wolffish	19.1	17.1	17.1	0.0					1.3	0.7

Values in metric tons of live weight

Sector and common pool include estimate of missing dealer reports

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting; (4) Observers and at-sea monitors via the Northeast Fisheries Observer Program. Differences with previous reports are due to corrections made to the database.

**Table 6: FY 2013 Northeast Multispecies Sector Carryover (mt)**

Stock	FY 2013 Available Annual Catch Entitlement (ACE)				FY 2013 Carryover	
	FY13 Initial ACE	FY12 Carryover	Total ACE	Total ACE as a Percent of Initial ACE	<i>de minimis</i>	Maximum
	A	B	A + B	C	D	E
GB Cod	1,776	447	2,223	125.2	17	93
GOM Cod	812	65	877	108.0	8	70
GB Haddock	26,111	2,695	28,807	110.3	170	917
GOM Haddock	185	64	249	134.4	2	18
GB Yellowtail Flounder	153	NA	152.6	100.0	NA	NA
SNE/MA Yellowtail Flounder	487	58.3	545.8	112.0	5	35
CC/GOM Yellowtail Flounder	466	100	565	121.4	5	25
Plaice	1,395	315	1,710	122.6	13	73
Witch Flounder	599	6	605	101.0	1	2
GB Winter Flounder	3,506	330	3,836	109.4	34	105
GOM Winter Flounder	688	68	756	109.8	7	38
SNE Winter Flounder	1,074	0	1,074	100.0	11	64
Redfish	10,092	822	10,914	108.1	104	556
White Hake	3,822	320	4,142	108.4	42	225
Pollock	12,801	1,230	14,031	109.6	129	696
Northern Windowpane	NA	NA	NA	NA	NA	NA
Southern Windowpane	NA	NA	NA	NA	NA	NA
Ocean Pout	NA	NA	NA	NA	NA	NA
Halibut	NA	NA	NA	NA	NA	NA
Wolffish	NA	NA	NA	NA	NA	NA

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting; (4) Observers and at-sea monitors via the Northeast Fisheries Observer Program. Differences with previous reports are due to corrections made to the database.

**Table 7: FY 2013 Northeast Multispecies Estimated State Water Sub-Component Catch Detail (mt)**

Stock	Total			Commercial			Recreational		
	Catch	Landings	Discard	Total Catch	Landings <sup>1</sup>	Discard <sup>1</sup>	Total Catch	Landings	Discard
	A+B+C+D	A+C	B+D	A+B	A	B	C+D	C	D
GB Cod	9.2	6.0	3.2	9.2	6.0	3.2	-	-	-
GOM cod	35.8	33.2	2.6	35.8	33.2	2.6	-*	-*	-*
GB Haddock	6.1	0.2	5.9	6.1	0.2	5.9	-	-	-
GOM Haddock	1.3	0.6	0.7	1.3	0.6	0.7	-*	-*	-*
GB Yellowtail Flounder	0.0	0.0	0.0	-	-	-	-	-	-
SNE/MA Yellowtail Flounder	14.5	8.3	6.1	14.5	8.3	6.1			
CC/GOM Yellowtail Flounder	42.8	40.8	2.0	42.8	40.8	2.0			
Plaice	19.6	10.4	9.2	19.6	10.4	9.2			
Witch Flounder	27.1	24.7	2.4	27.1	24.7	2.4			
GB Winter Flounder	0.0	0.0	0.0	-	-	-	-	-	-
GOM Winter Flounder	67.4	65.3	2.1	37.1	37.0	0.1	30.3	28.3	2.0
SNE/MA Winter Flounder	55.7	47.7	8.1	48.0	44.0	4.0	7.7	3.7	4.0
Redfish	19.0	6.5	12.5	19.0	6.5	12.5			
White Hake	2.3	0.8	1.5	2.3	0.8	1.5			
Pollock	981.7	258.2	723.6	8.6	3.9	4.7	973.1	254.2	718.9
Northern Windowpane	0.9	0.0	0.9	0.9	-	0.9			
Southern Windowpane	37.3	20.8	16.5	37.3	20.8	16.5			
Ocean Pout	1.5	0.0	1.5	1.5	-	1.5			
Halibut	22.8	21.1	1.7	22.8	21.1	1.7			
Wolffish	1.3	0.0	1.3	1.3	0.0	1.3			

\*Recreational catch of GOM cod and haddock in state waters is attributed to the recreational sub-ACL (see Tables 1 - 5), and so is not counted above.

<sup>1</sup>January through April 2014 commercial catches are estimated.

State discard rate estimates based on discard rates on federal trips

Values in metric tons of live weight

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting; (4) Observers and at-sea monitors via the Northeast Fisheries Observer Program. Differences with previous reports are due to corrections made to the database.

**Table 8: FY 2013 Northeast Multispecies Other Sub-Component Catch Detail (mt)**

Stock	Total Catch	SCALLOP <sup>1</sup>	FLUKE	HAGFISH	HERRING	LOBSTER/ CRAB	MENHADEN	MONKFISH	RESEARCH	SCUP	SHRIMP
GB cod	34.2	4.9	0.3	0.0	1.4	0.8	0.3	0.2	14.5	0.1	0.0
GOM cod	2.9	0.2	-	0.0	1.3	0.3	-	-	0.1	-	-
GB Haddock	56.5	3.5	0.1	0.0	5.2*	0.0	0.0	0.0	0.5	0.1	0.0
GOM Haddock	1.6	0.0	-	-	0.3*	-	-	-	0.0	-	-
GB Yellowtail Flounder	0.0	-*	-	-	-*	-	-	-	-	0.0	-
SNE Yellowtail Flounder	29.8	-*	5.7	-	1.3	0.0	0.0	0.0	1.3	5.6	0.0
CC/GOM Yellowtail Flounder	29.7	23.2	-	-	1.3	-	-	-	2.1	-	-
Plaice	29.8	13.5	0.7	-	1.3	0.0	0.0	0.0	0.8	0.8	0.0
Witch Flounder	75.8	26.7	5.7	0.0	3.3	0.1	0.1	0.0	0.6	4.7	0.0
GB Winter Flounder	41.0	25.0	-	-	1.5	-	-	-	-	0.1	-
GOM Winter Flounder	8.9	6.0	-	0.0	0.2	0.0	-	-	0.1	-	-
SNE Winter Flounder	181.6	78.2	10.8	-	4.7	0.0	0.1	0.0	19.9	9.7	0.0
Redfish	3.9	0.0	0.0	0.0	1.0	0.4	0.1	0.0	0.0	0.0	0.0
White Hake	8.3	1.0	0.1	0.0	2.0	1.0	0.3	0.1	0.1	0.2	0.0
Pollock	1,132.4	0.0	0.0	0.0	0.6	0.1	0.0	0.0	0.1	0.0	0.0
Northern Windowpane	41.6	40.7	-	0.0	0.2	0.0	-	-	0.0	0.0	-
Southern Windowpane	272.4	-*	66.9	-	3.0	0.1	0.5	0.0	0.0	69.6	0.0
Ocean Pout	24.6	2.9	0.5	0.0	2.0	0.0	0.0	0.0	0.0	0.5	0.0
Halibut <sup>2</sup>	1.5	0.2	0.0	0.0	0.1	0.6	0.0	0.0	0.0	0.0	0.0
Wolfish	0.7	0.5	0.0	-	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Values in metric tons of live weight

<sup>1</sup>Based on scallop fishing year March, 2013 through February, 2014

<sup>2</sup> Note some Canadian landings of this stock are included in the most recent assessment for Atlantic halibut (2012 Assessment Update). However, Canadian landings for 2013 have not yet been reported to the Northwest Atlantic Fisheries Organization (NAFO), and, as a result, are not included here.

\*Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not included above.

Source: NMFS Greater Atlantic Regional Office  
October 29, 2014, run date of July 15, 2014

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 8: FY 2013 Northeast Multispecies Other Sub-Component Catch Detail (mt)**

Stock	Total Catch	SQUID	SQUID/ WHITING	SURFCLAM	TILEFISH	WHELK/ CONCH	WHITING	UNKNOWN	RECREATIONAL
GB cod	34.2	0.6	0.6	0.0	0.0	0.1	0.0	2.5	8.0
GOM cod	2.9	-	0.2	-	-	0.0	0.4	0.5	-*
GB Haddock	56.5	14.8	15.5	0.0	0.0	0.0	0.0	16.7	-
GOM Haddock	1.6	-	0.3	-	-	-	0.5	0.5	-*
GB Yellowtail Flounder	0.0	-*	0.0*	-	-	-	-	0.0*	
SNE Yellowtail Flounder	29.8	2.0	2.2	-	-	-	0.0	11.7	
CC/GOM Yellowtail Flounder	29.7	-	0.6	-	-	-	1.3	1.3	
Plaice	29.8	3.6	3.9	-	-	-	0.1	5.1	
Witch Flounder	75.8	8.7	9.9	0.0	0.0	0.0	0.2	15.8	
GB Winter Flounder	41.0	0.5	12.7	-	-	-	-	1.3	
GOM Winter Flounder	8.9	-	0.1	-	-	0.0	0.2	2.3	0.1
SNE Winter Flounder	181.6	14.5	11.2	-	-	-	0.0	32.4	0.0
Redfish	3.9	0.6	0.7	0.0	0.0	0.0	0.0	0.9	
White Hake	8.3	0.6	1.3	0.0	0.0	0.1	0.0	1.5	
Pollock	1,132.4	1.0	1.0	0.0	0.0	0.0	0.0	1.3	1,128.0
Northern Windowpane	41.6	0.0	0.6	0.0	-	0.0	0.0	0.1	
Southern Windowpane	272.4	12.3	19.0	0.0	0.0	0.0	0.0	100.9	
Ocean Pout	24.6	5.6	5.9	0.0	0.0	0.0	0.1	6.9	
Halibut <sup>2</sup>	1.5	0.1	0.2	0.0	0.0	0.0	0.0	0.3	
Wolfish	0.7	0.0	0.0	-	-	-	0.0	0.1	

Values in metric tons of live weight

<sup>1</sup>Based on scallop fishing year March, 2013 through February, 2014

<sup>2</sup> Note some Canadian landings of this stock are included in the most recent assessment for Atlantic halibut (2012 Assessment Update). However, Canadian landings for 2013 have not yet been reported to the Northwest Atlantic Fisheries Organization (NAFO), and, as a result, are not included here.

\*Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not included above.

Source: NMFS Greater Atlantic Regional Office  
October 29, 2014, run date of July 15, 2014

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 9: FY 2013 Northeast Multispecies Other Sub-Component Landings Detail (mt)**

Stock	Total	SCALLOP <sup>1</sup>	FLUKE	HAGFISH	HERRING	LOBSTER/ CRAB	MENHADEN	MONKFISH	RESEARCH	SCUP	SHRIMP
GB cod	22.7	0.2	0.2	-	0.1	0.1	-	0.2	14.5	0.0	-
GOM cod	0.2	-	-	-	-	0.1	-	-	0.1	-	-
GB Haddock	0.5	-	0.0	-	-*	0.0	-	-	0.5	-	-
GOM Haddock	0.0	-	-	-	-*	-	-	-	0.0	-	-
GB Yellowtail Flounder	-	-*	-	-	-	-	-	-	-	-	-
SNE Yellowtail Flounder	5.3	-*	0.4	-	0.7	0.0	-	-	1.3	0.1	-
CC/GOM Yellowtail Flounder	2.4	0.3	-	-	-	-	-	-	2.1	-	-
Plaice	0.8	0.0	0.0	-	-	-	-	-	0.8	-	-
Witch Flounder	2.2	0.2	1.3	-	-	-	-	-	0.6	0.0	-
GB Winter Flounder	0.7	0.7	-	-	-	-	-	-	-	-	-
GOM Winter Flounder	2.2	-	-	-	-	-	-	-	0.1	-	-
SNE Winter Flounder	29.2	0.6	2.0	-	0.4	0.0	-	0.0	19.9	0.4	-
Redfish	0.6	-	-	-	0.3	0.1	-	-	-	-	-
White Hake	1.3	-	0.0	-	-	0.1	-	0.0	0.1	0.2	-
Pollock	450.2	-	0.0	-	-	0.0	-	-	0.1	0.0	-
Northern Windowpane	-	-	-	-	-	-	-	-	-	-	-
Southern Windowpane	2.0	-*	0.8	-	-	-	-	-	-	-	-
Ocean Pout	0.0	-	-	-	-	0.0	-	-	-	-	-
Halibut <sup>2</sup>	0.8	0.0	-	-	-	0.6	-	0.0	0.0	-	-
Wolfish	-	-	-	-	-	-	-	-	-	-	-

Values in metric tons of live weight

<sup>1</sup>Based on scallop fishing year March, 2013 through February, 2014

<sup>2</sup> Note some Canadian landings of this stock are included in the most recent assessment for Atlantic halibut (2012 Assessment Update). However, Canadian landings for 2013 have not yet been reported to the Northwest Atlantic Fisheries Organization (NAFO), and, as a result, are not included here.

\*Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not included above.

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 9: FY 2013 Northeast Multispecies Other Sub-Component Landings Detail (mt)**

Stock	Total	SQUID	SQUID/ WHITING	SURFCLAM	TILEFISH	WHELK/ CONCH	WHITING	UNKNOWN	RECREATIONAL
GB cod	22.7	0.0	0.0	-	-	-	-	1.3	6.1
GOM cod	0.2	-	-	-	-	-	-	0.1	-*
GB Haddock	0.5	-	-	-	-	-	-	-	0.0
GOM Haddock	0.0	-	-	-	-	-	-	0.0	-*
GB Yellowtail Flounder	-	-	-	-	-	-	-	-	-
SNE Yellowtail Flounder	5.3	0.0	-	-	-	-	-	2.8	-
CC/GOM Yellowtail Flounder	2.4	-	-	-	-	-	-	0.0	-
Plaice	0.8	-	-	-	-	-	-	0.0	-
Witch Flounder	2.2	-	0.1	-	-	-	-	0.1	-
GB Winter Flounder	0.7	-	-	-	-	-	-	-	-
GOM Winter Flounder	2.2	-	-	-	-	-	-	2.0	0.1
SNE Winter Flounder	29.2	0.3	0.1	-	-	-	0.0	5.4	-
Redfish	0.6	0.0	0.1	-	-	-	0.0	0.0	-
White Hake	1.3	-	0.7	-	-	0.0	-	0.2	-
Pollock	450.2	0.0	0.0	-	-	-	-	0.1	449.9
Northern Windowpane	-	-	-	-	-	-	-	-	-
Southern Windowpane	2.0	-	-	-	-	-	0.0	1.2	-
Ocean Pout	0.0	-	-	-	-	-	-	0.0	-
Halibut <sup>2</sup>	0.8	-	-	-	-	-	-	0.2	-
Wolffish	-	-	-	-	-	-	-	-	-

Values in metric tons of live weight

<sup>1</sup>Based on scallop fishing year March, 2013 through February, 2014

<sup>2</sup> Note some Canadian landings of this stock are included in the most recent assessment for Atlantic halibut (2012 Assessment Update). However, Canadian landings for 2013 have not yet been reported to the Northwest Atlantic Fisheries Organization (NAFO), and, as a result, are not included here.

\*Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not included above.

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 10: FY 2013 Northeast Multispecies Other Sub-Component Estimated Discards Detail (mt)**

Stock	Total	SCALLOP <sup>1</sup>	FLUKE	HAGFISH	HERRING	LOBSTER/ CRAB	MENHADEN	MONKFISH	RESEARCH	SCUP	SHRIMP
GB cod	11.5	4.7	0.1	0.0	1.3	0.7	0.3	0.1	0.0	0.1	0.0
GOM cod	2.7	0.2	-	0.0	1.3	0.2	-	-	0.0	-	-
GB Haddock	56.0	3.5	0.1	0.0	5.2*	0.0	0.0	0.0	0.0	0.1	0.0
GOM Haddock	1.6	0.0	-	-	.3*	-	-	-	-	-	-
GB Yellowtail Flounder	0.0	-*	-	-	-*	-	-	-	-	0.0	-
SNE Yellowtail Flounder	24.5	-*	5.3	-	0.5	0.0	0.0	0.0	0.0	5.5	0.0
CC/GOM Yellowtail Flounder	27.3	22.8	-	-	1.3	-	-	-	-	-	-
Plaice	28.9	13.5	0.7	-	1.3	0.0	0.0	0.0	0.0	0.8	0.0
Witch Flounder	73.6	26.5	4.4	0.0	3.3	0.1	0.1	0.0	0.0	4.7	0.0
GB Winter Flounder	40.3	24.2	-	-	1.5	-	-	-	-	0.1	-
GOM Winter Flounder	6.7	6.0	-	0.0	0.2	0.0	-	-	0.0	-	-
SNE Winter Flounder	152.4	77.6	8.8	-	4.3	0.0	0.1	0.0	0.0	9.3	0.0
Redfish	3.3	0.0	0.0	0.0	0.8	0.3	0.1	0.0	0.0	0.0	0.0
White Hake	7.1	1.0	0.1	0.0	2.0	0.9	0.3	0.1	0.0	0.1	0.0
Pollock	682.2	0.0	0.0	0.0	0.6	0.1	0.0	0.0	0.0	0.0	0.0
Northern Windowpane	41.6	40.7	-	0.0	0.2	0.0	-	-	0.0	0.0	-
Southern Windowpane	270.4	-*	66.1	-	3.0	0.1	0.5	0.0	0.0	69.6	0.0
Ocean Pout	24.6	2.9	0.5	0.0	2.0	0.0	0.0	0.0	0.0	0.5	0.0
Halibut	0.7	0.2	0.0	0.0	0.1	0.0	0.0	0.0	0.0	0.0	0.0
Wolfish	0.7	0.5	0.0	-	0.0	0.0	0.0	0.0	0.0	0.0	0.0

Values in metric tons of live weight

<sup>1</sup>Based on scallop fishing year March, 2013 through February, 2014

\*Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not included above.

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 10: FY 2013 Northeast Multispecies Other Sub-Component Estimated Discards Detail (mt)**

Stock	Total	SQUID	SQUID/ WHITING	SURFCLAM	TILEFISH	WHELK/ CONCH	WHITING	UNKNOWN	RECREATIONAL
GB cod	11.5	0.6	0.6	0.0	0.0	0.1	0.0	1.2	1.9
GOM cod	2.7	-	0.2	-	-	0.0	0.4	0.4	-*
GB Haddock	56.0	14.8	15.5	0.0	0.0	0.0	0.0	16.7	-
GOM Haddock	1.6	-	0.3	-	-	-	0.5	0.5	-*
GB Yellowtail Flounder	0.0	-*	0.0*	-	-	-	-	0.0*	
SNE Yellowtail Flounder	24.5	1.9	2.2	-	-	-	0.0	9.0	
CC/GOM Yellowtail Flounder	27.3	-	0.6	-	-	-	1.3	1.3	
Plaice	28.9	3.6	3.9	-	-	-	0.1	5.0	
Witch Flounder	73.6	8.7	9.8	0.0	0.0	0.0	0.2	15.8	
GB Winter Flounder	40.3	0.5	12.7	-	-	-	-	1.3	
GOM Winter Flounder	6.7	-	0.1	-	-	0.0	0.2	0.2	0.0
SNE Winter Flounder	152.4	14.2	11.1	-	-	-	0.0	27.0	0.0
Redfish	3.3	0.6	0.6	0.0	0.0	0.0	0.0	0.9	
White Hake	7.1	0.6	0.6	0.0	0.0	0.1	0.0	1.3	
Pollock	682.2	1.0	1.0	0.0	0.0	0.0	0.0	1.2	678.1
Northern Windowpane	41.6	0.0	0.6	0.0	-	0.0	0.0	0.1	
Southern Windowpane	270.4	12.3	19.0	0.0	0.0	0.0	0.0	99.7	
Ocean Pout	24.6	5.6	5.9	0.0	0.0	0.0	0.1	6.9	
Halibut	0.7	0.1	0.2	0.0	0.0	0.0	0.0	0.2	
Wolfish	0.7	0.0	0.0	-	-	-	0.0	0.1	

Values in metric tons of live weight

<sup>1</sup>Based on scallop fishing year March, 2013 through February, 2014

\*Some or all catch attributed to separate sub-ACL as shown in Tables 1 through 5, and so is not included above.

Source: NMFS Greater Atlantic Regional Office

October 29, 2014, run date of July 15, 2014

These criteria are used by the Northeast Regional Office (NERO) to categorize trips to attribute groundfish catch for groundfish ACL accounting. By necessity these rules cannot capture the full complexity of categorizing every trip taken by vessels fishing in the Northeast. Further analysis should be completed to definitively attribute groundfish catch to an FMP for management purposes.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 11: FY 2013 U.S. Catch of Transboundary U.S./Canada Stocks (mt)**

Stock	Total U.S. Catch	U.S. Catch by Fishery Component								
		Groundfish	Sector	Common Pool	Recreational	Herring Fishery	Scallop Fishery	Small Mesh Fisheries	State Water	Other
		A to H	A+B+C	A	B	C	D	E	F	G
Eastern GB Cod	33.6	33.3	33.3	-						0.3
Eastern GB Haddock	638.9	578.8	578.8	-		58.0				2.0
GB Yellowtail Flounder	93.3	55.8	55.8	0.0			37.5	2.5		0.0

\*Landings extrapolated from observer data.  
 Values in live weight  
 Includes estimate of missing dealer reports  
 October 29, 2014, run date of July 15, 2014

**Table 12: FY 2013 U.S./Canada Groundfish Trips, Vessels, and A DAS Used**

Area <sup>1</sup>	Number of Vessels		Number of Trips		A DAS Used		Number of Observed Trips	
	Sector	Common Pool	Sector	Common Pool	Sector	Common Pool	Sector	Common Pool
Eastern U.S./Canada Area	55	0	305	0	2,155	0	36	0
Western U.S./Canada Area	67	1	731	4	4,763	25	162	0
Total	70	1	766	4	4,976	25	164	0

<sup>1</sup>Area based on area fished. Totals don't sum due to multi-area trips

Source: NMFS Greater Atlantic Regional Fisheries Office  
 October 29, 2014, run date of July 15, 2014

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 11A: FY 2013 U.S. Landings of Transboundary U.S./Canada Stocks (mt)**

Stock	U.S. Landings	U.S. Catch by Fishery Component								
		Groundfish	Sector	Common Pool	Recreational	Herring Fishery	Scallop Fishery	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
Eastern GB Cod	23.4	23.4	23.4	-						-
Eastern GB Haddock	582.2	528.7	528.7	-		53.5				-
GB Yellowtail Flounder	49.7	46.2	46.2	0.0			3.6	-		-

Values in live weight

Includes estimate of missing dealer reports

Source: NMFS Greater Atlantic Regional Fisheries Office  
October 29, 2014, run date of July 15, 2014

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 11B: FY 2013 Estimated U.S. Discards of Transboundary U.S./Canada Stocks (mt)**

Stock	U.S. Discards	U.S. Catch by Fishery Component								
		Groundfish	Sector	Common Pool	Recreational	Herring Fishery	Scallop Fishery	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
Eastern GB Cod	10.2	9.9	9.9	-						0.3
Eastern GB Haddock	56.7	50.2	50.2	-		4.5				2.0
GB Yellowtail Flounder	43.5	9.6	9.6	0.0			33.9	2.5		0.0

Values in live weight

Includes estimate of missing dealer reports

Source: NMFS Greater Atlantic Regional Fisheries Office

October 29, 2014, run date of July 15, 2014

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 13: FY 2013 U.S. TACs of Transboundary U.S./Canada Stocks (mt)**

Stock	U.S. TAC	Fishery Component Allocation								
		Groundfish	Sector*	Common Pool	Recreational	Herring Fishery	Scallop Fishery	Small-Mesh Fisheries	State Water	Other
		A+B+C	A	B	C	D	E	F	G	H
Eastern Georges Bank Cod	96	92	90	2						
Eastern Georges Bank Haddock	3,952	3,754	3,742	12						
Georges Bank Yellowtail Flounder	215.0	154.5	152.6	1.9			41.5	4.0		4.3

\*Sector sub-ACL values have been corrected from the table dated September 23, 2014 to include unrenewed permit potential sector contributions

Values in live weight

Source: NMFS Greater Atlantic Regional Fisheries Office  
October 29, 2014

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 14: FY 2013 Percent of U.S. TAC Caught for Transboundary U.S./Canada Stocks (%)**

Stock	% of U.S. TAC	Percent of Each Fishery Component U.S. TAC Caught								
		Groundfish	Sector*	Common Pool	Recreational	Herring Fishery	Scallop Fishery	Small Mesh Fisheries	State Water	Other
	A to H	A+B+C	A	B	C	D	E	F	G	H
Eastern GB Cod	35.0	36.2	37.0	-						NA
Eastern GB Haddock	16.2	15.4	15.5	-		NA				NA
GB Yellowtail Flounder	43.4	36.1	36.5	0.4			90.3	63.7		0.6

\*Sector sub-ACL values have been corrected from the report dated October 20, 2014 to include unrenewed permit potential sector contributions

Values in percent live weight (%)

Includes estimate of missing dealer reports

Source: NMFS Greater Atlantic Regional Fisheries Office  
October 29, 2014, run date of July 15, 2014

Any value for a non-allocated species may be due to landings of that stock; misreporting of species and/or stock area; and/or estimated landings (in lieu of missing reports) based on vessel histories.

These data are the best available to NOAA's National Marine Fisheries Service (NMFS). Data sources for this report include: (1) Vessels via VMS; (2) Vessels via vessel logbook reports; (3) Dealers via Dealer Electronic reporting. Differences with previous reports are due to corrections made to the database.

**Table 15**  
**U.S./Canada**  
**Commercial**  
**Groundfish**  
 FY13-FY14 to  
 Date

STOCK	Fishing_Year	Month	Sum of KEPT_MT	Sum of DISCARD_MT	Sum of CATCH_MT	
GB Cod East	2013	May	0.9	1.9	2.8	
	2013	June	0.9	4.9	5.8	
	2013	July	1.6	0.6	2.1	
	2013	August	0.6	0.2	0.8	
	2013	September	0.3	0.1	0.4	
	2013	October	0.9	0.4	1.4	
	2013	November	0.9	0.2	1.1	
	2013	December	1.3	0.3	1.6	
	2013	January	0.2	0.1	0.3	
	2013	February	1.1	0.1	1.2	
	2013	March	1.9	0.6	2.5	
	2013	April	12.8	0.5	13.3	
	<b>2013</b>	<b>Total</b>	<b>23.4</b>	<b>9.9</b>	<b>33.3</b>	
	2014	May	3.3	0.6	4.0	
	2014	June	1.8	0.4	2.2	
	2014	July	2.3	0.1	2.4	
	2014	August	2.8	0.1	2.9	
	<b>2014</b>	<b>Total</b>	<b>10.3</b>	<b>1.3</b>	<b>11.5</b>	
	<b>GB Cod East Total</b>			<b>33.6</b>	<b>11.2</b>	<b>44.8</b>
	GB Haddock East	2013	May	6.3	5.3	11.7
2013		June	9.0	13.8	22.7	
2013		July	33.0	4.5	37.5	
2013		August	22.4	2.3	24.7	
2013		September	25.2	2.2	27.5	
2013		October	32.1	4.5	36.6	
2013		November	24.2	2.1	26.2	
2013		December	85.6	4.0	89.7	
2013		January	22.2	1.7	23.8	
2013		February	30.5	1.6	32.1	
2013		March	157.1	4.3	161.4	
2013		April	81.1	3.8	84.9	
<b>2013</b>		<b>Total</b>	<b>528.7</b>	<b>50.2</b>	<b>578.8</b>	
2014		May	201.9	19.2	221.1	
2014		June	148.2	9.3	157.6	
2014		July	24.1	1.6	25.7	
2014		August	24.0	4.3	28.3	
<b>2014</b>		<b>Total</b>	<b>398.3</b>	<b>34.5</b>	<b>432.7</b>	
<b>GB Haddock East Total</b>				<b>927.0</b>	<b>84.6</b>	<b>1,011.6</b>
GB Yellowtail Flounder		2013	May	3.2	0.5	3.7
	2013	June	1.8	0.4	2.2	
	2013	July	2.7	1.4	4.1	
	2013	August	3.5	1.3	4.8	
	2013	September	1.3	0.7	1.9	
	2013	October	1.3	1.2	2.5	
	2013	November	1.6	1.1	2.6	
	2013	December	4.3	0.7	5.1	
	2013	January	0.2	0.4	0.6	
	2013	February	0.1	0.7	0.8	
	2013	March	0.2	0.4	0.6	
	2013	April	26.1	0.8	26.9	
	<b>2013</b>	<b>Total</b>	<b>46.2</b>	<b>9.6</b>	<b>55.8</b>	
	2014	May	9.0	1.5	10.5	
	2014	June	7.2	2.0	9.2	
	2014	July	2.3	0.9	3.2	
2014	August	2.9	0.3	3.3		
<b>2014</b>	<b>Total</b>	<b>21.5</b>	<b>4.7</b>	<b>26.2</b>		
<b>GB Yellowtail Flounder Total</b>			<b>67.6</b>	<b>14.3</b>	<b>81.9</b>	

Through Aug. 2014  
 Groundfish trips only  
 Live Weight  
 Groundfish landings  
 include estimate of  
 missing dealer  
 reports  
 Source: NMFS  
 Greater Atlantic  
 Regional Office  
 September 8, 2014

**Table 16: FY 2011 - 2013 GOM Cod and Haddock Recreational Catch Evaluation  
(mt)**

Stock	Fishing Year	Recreational Catch				
		Catch	Landings	Discard	Recreational sub-ACL	Percent of Catch Limit Taken
		A + B	A	B		
GOM Cod	2011	1,640.3	1,640.3	NA*	2,824	58.1
	2012	937.4	634.6	302.8	2,215	42.3
	2013	639.3	540.6	98.8	486	131.5
	<b>Average</b>	<b>1,072.4</b>	<b>938.5</b>	<b>133.8</b>	<b>1,842</b>	<b>58.2</b>
GOM Haddock	2011	238.5	238.5	NA*	308	77.4
	2012	280.7	280.7	NA*	259	108.4
	2013	231.5	231.5	0.0	74	312.2
	<b>Average</b>	<b>250.2</b>	<b>250.2</b>	<b>0.0</b>	<b>214</b>	<b>117.1</b>

\*Estimates not applicable. Recreational discards are not attributed to the ACL consistent with the most recent assessments for these stocks used to set the respective quotas (GARM III). SARC 53 for GOM cod included recreational discards for the first time and was used to set quotas beginning in FY 2012. SARC 59 for GOM haddock included recreational discards for the first time and will be used to set quotas starting in FY 2015.

Recreational estimates based on Marine Recreational Information Program (MRIP) data.  
Values in metric tons of live weight

Source: NMFS Greater Atlantic Regional Office  
October 29, 2014

These data are the best available to NOAA's National Marine Fisheries Service (NMFS).